



Geochemistry of PM₁₀ over Europe during the EMEP intensive measurement periods in summer 2012 and winter 2013

Andrés Alastuey¹, Xavier Querol¹, Wenche Aas², Franco Lucarelli³, Noemí Pérez¹, Teresa Moreno¹, Fabrizia Cavalli⁴, Hans Areskoug⁵, Violeta Balan⁶, Maria Catrambone⁷, Darius Ceburnis⁸, José C. Cerro⁹, Sébastien Conil¹⁰, Lusine Gevorgyan¹¹, Christoph Hueglin¹², Kornelia Imre¹³, Jean-Luc Jaffrezo¹⁴, Sarah R. Leeson¹⁵, Nikolaos Mihalopoulos^{16,24}, Marta Mitosinkova¹⁷, Colin D. O'Dowd⁸, Jorge Pey¹⁸, Jean-Philippe Putaud⁶, Véronique Riffault¹⁹, Anna Ripoll¹, Jean Sciare^{20,21}, Karine Sellegri²², Gerald Spindler²³, and Karl Espen Yttri²

¹Department of Geosciences, Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona, Spain

²Atmosphere and Climate Department, NILU-Norwegian Institute for Air Research, Kjeller, Norway

³Dipartimento di Fisica e Astronomia and National Institute of Nuclear Physics (INFN), Sesto Fiorentino, Florence, Italy

⁴European Commission – DG Joint Research Centre, Ispra, Italy

⁵Department of Environmental Sciences and Analytical Chemistry, Stockholm University, ACES, Stockholm, Sweden

⁶Hydrometeorologic State Service, Ministry of Ecology and Natural Resources, Chisinau, Moldova

⁷CNR, Institute of Atmospheric Pollution Research, Monterotondo Stazione, Rome, Italy

⁸School of Physics, National University of Ireland Galway, Galway, Ireland

⁹Laboratory of Environmental Analytical Chemistry, Illes Balears University, Palma de Mallorca, Spain

¹⁰ANDRA – DRD – Observation Surveillance, Observatoire Pérenne de l'Environnement, Bure, France

¹¹Environmental Impact Monitoring Center, Yerevan, Armenia

¹²Empa, Swiss Federal Laboratories for Materials Science and Technology, 8600 Dübendorf, Switzerland

¹³MTA-PE Air Chemistry Research Group, University of Veszprém, Veszprém, Hungary

¹⁴Laboratoire de Glaciologie et Géophysique de l'Environnement, UGA-CNRS, St. Martin d'Hères CEDEX, France

¹⁵Centre for Ecology and Hydrology (CEH), Bush Estate, Penicuik, EH26 0QB, UK

¹⁶Environmental Chemical Processes Laboratory, University of Crete, Heraklion, Greece

¹⁷Department of Air Quality, Slovak Hydrometeorological Institute, Bratislava, Slovak Republic

¹⁸Spanish Geological Survey, Zaragoza IGME Unit, Zaragoza, Spain

¹⁹Département Sciences de l'Atmosphère et Génie de l'Environnement (SAGE), Mines Douai, Douai, France

²⁰Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

²¹The Cyprus Institute, Energy, Environment and Water Research Center, Nicosia, Cyprus

²²Laboratoire de Météorologie Physique LaMP-CNRS/OPGC, Aubière, France

²³Department of Atmospheric Chemistry, Leibniz Institute for Tropospheric Research (TROPOS) Leipzig, Germany

²⁴Institute for Environmental Research and Sustainable Development, National Observatory of Athens, Pendeli, Greece

Correspondence to: Andrés Alastuey (andres.alastuey@idaea.csic.es)

Received: 15 January 2016 – Published in Atmos. Chem. Phys. Discuss.: 24 February 2016

Revised: 25 April 2016 – Accepted: 26 April 2016 – Published: 19 May 2016

Abstract. The third intensive measurement period (IMP) organised by the European Monitoring and Evaluation Programme (EMEP) under the UNECE CLTRAP took place in summer 2012 and winter 2013, with PM₁₀ filter samples concurrently collected at 20 (16 EMEP) regional background sites across Europe for subsequent analysis of their mineral dust content. All samples were analysed by the same or a comparable methodology. Higher PM₁₀ mineral dust loadings were observed at most sites in summer ($0.5\text{--}10\ \mu\text{g m}^{-3}$) compared to winter ($0.2\text{--}2\ \mu\text{g m}^{-3}$), with the most elevated concentrations in the southern- and easternmost countries, accounting for 20–40 % of PM₁₀. Saharan dust outbreaks were responsible for the high summer dust loadings at western and central European sites, whereas regional or local sources explained the elevated concentrations observed at eastern sites. The eastern Mediterranean sites experienced elevated levels due to African dust outbreaks during both summer and winter. The mineral dust composition varied more in winter than in summer, with a higher relative contribution of anthropogenic dust during the former period. A relatively high contribution of K from non-mineral and non-sea-salt sources, such as biomass burning, was evident in winter at some of the central and eastern European sites. The spatial distribution of some components and metals reveals the influence of specific anthropogenic sources on a regional scale: shipping emissions (V, Ni, and SO₄²⁻) in the Mediterranean region, metallurgy (Cr, Ni, and Mn) in central and eastern Europe, high temperature processes (As, Pb, and SO₄²⁻) in eastern countries, and traffic (Cu) at sites affected by emissions from nearby cities.

1 Introduction

Mineral dust along with sea-salt aerosols are the major components of particulate matter (PM) mass in the atmosphere on a global scale (IPCC, 2007), playing a key role in the planet (Knippertz and Stuut, 2014). Mineral dust particles are generated mainly by wind erosion and soil resuspension in deserts and arid regions (e.g. Zhao et al., 2010; Kok, 2011), and their size distribution is characterised by a coarse size mode with a small fraction in the accumulation mode. Atmospheric residence time varies from less than 1 day to more than 1 week, depending on particle size and composition, but essentially on the effect of mesoscale and synoptic meteorology.

The magnitude of dust emissions to the atmosphere depends on the surface wind speed and soil-related factors such as texture, moisture and vegetation cover. Currently there is still a high uncertainty in the estimates of global dust emissions varying from 500 to 3000 Mt year⁻¹ according to models (Huneus et al., 2011). The major mineral dust sources from arid and semi-arid continental regions are located in subtropical areas in the Northern Hemisphere, and

the Sahara–Sahel–Chad dust corridor is considered as the largest one of these (Prospero et al., 2002; Engelstaedter et al., 2006; Moreno et al., 2006). The proximity of southern Europe to these large dust emitting areas and region-specific meteorological dynamics are responsible for the frequent transport of dust over southern Europe and the subsequent deterioration of air quality (e.g. Bergametti et al., 1989; Querol et al., 1998, 2009; Rodríguez et al., 2001; Kallos et al., 2006; Mitsakou et al., 2008; Pey et al., 2013).

Natural mineral dust mainly consists of silicate, carbonate, phosphate and oxide/hydroxide minerals derived from the erosion and weathering of rocks and soils (Moreno et al., 2008; Scheuven et al., 2013). Mineralogical characterisation of desert dust aerosols in northern Africa carried out during the SAMUM campaign showed that the major mineral constituents of the aerosol were quartz, potassium feldspar, plagioclase, calcite, hematite and the clay minerals illite, kaolinite and chlorite (Kandler et al., 2009; Scheuven et al., 2011). The chemical composition of airborne mineral dust depends on the geology of the source region, reflecting the composition of the parent soil of the source area. However, changes in chemical composition and physical characteristics of mineral dust may take place during transport due to the preferential deposition of coarse particles (Aluko and Noll, 2006; Scheuven et al., 2013) and to the interaction with other particulate and gaseous pollutants (Rodríguez et al., 2011), resulting in coating or mixing of dust particles (Kandler et al., 2007; Levin and Ganor, 1996). However, these interactions are not always large and the chemical modifications can be minor (Aymoz et al., 2004).

Mineral dust particles are also emitted by anthropogenic sources, such as agricultural activities, construction sites, mining, certain industrial activities such as the cement and ceramic industries, and road dust resuspension (Zender et al., 2004). However, there is a much variation in the contribution of such emissions depending on geographic location (Tegen et al., 2004). Whereas recent satellite observations suggest that the anthropogenic fraction of suspended mineral dust on the global scale represents only 20 to 25 % of the total (Ginoux et al., 2012), on local or regional scales the anthropogenic contribution can rise to become the dominant source and so highly affect air quality (Querol et al., 2004).

Research on mineral dust is of great interest due to the impact of such particles on air quality, health and climate. Inhalation of mineral dust has potential adverse effects on health caused by particle composition, shape and particle size (Morman and Plumlee, 2014). Numerous scientific publications have demonstrated a relationship between the increase in PM₁₀ concentration during Saharan dust events and severe health outcomes in the southern European region (Pérez et al., 2008, 2012; Diaz et al., 2012; Mallone et al., 2011; Stafoggia et al., 2013). For instance, Pérez et al. (2008, 2012) evidenced that short-term exposure to PM during Saharan dust days is associated with both cardiovascular and respiratory mortality in Barcelona. However, as shown in the lit-

erature review by Karanasiou et al. (2012), the studies published show contradictory results regarding the health impact of Saharan dust outbreaks. Thus, further studies on chemical characterisation, mixing state and potential toxicity of coarse particles transported from the Saharan desert are needed to clarify this issue.

Mineral dust can impact climate by different mechanisms, but mainly by interaction with radiative forcing and by modification of cloud properties (see references in Highwood and Ryder, 2014; Miller et al., 2014; and Nenes et al., 2014). Dust particles may scatter and/or absorb solar radiation and also act as ice nuclei. Net global radiative forcing of mineral dust is estimated as -0.1 (-0.3 to $+0.1$) W m^{-2} (IPCC, 2013). The optical properties of dust particles depend on its mineralogical and chemical composition, with hematite having the largest absorption of UV and visible spectra (Lafon et al., 2006) among the inorganic components.

PM mass concentrations are routinely measured in national and regional Air Quality networks, such as the European environmental agency's (EEA) air quality programme (EC, 2008) and the European Monitoring and Evaluation Programme (EMEP) (UNECE, 2009; Tørseth et al., 2012). Some PM components such as SO_4^{2-} , NO_3^- , NH_4^+ , organic carbon (OC) and elemental carbon (EC) are also measured at sites included in the referred networks.

Even though the measurement of mineral dust is a compulsory part of the EMEP monitoring program (UNECE, 2009), and chemical speciation is recommended by the World Meteorological Organisation (WMO) Global Atmospheric Watch (GAW) (WMO, 2003, 2007), it is not generally included as a routine part of the monitoring programmes, probably due to the lack of a reference method and because in some areas it has been considered as a minor component of PM. In some studies, mineral components are estimated from the analysis of the soluble fraction of one or few mineral elements (Rodríguez et al., 2012). This may result in the underestimation (or even overestimation) of the mineral load, given that most minerals are water insoluble.

The chemical composition of mineral matter has been the topic of several studies conducted across Europe (Querol et al., 2001; Alemón et al., 2004; Moreno et al., 2006; Lucarelli et al., 2011; Nava et al., 2012). Putaud et al. (2010) compiled data on physical and chemical characteristics of PM₁₀ and PM_{2.5}, including mineral dust, from 60 sites across Europe, but different sampling techniques and analytical methodologies hamper the comparability. A review on methods for long-term in situ characterisation of aerosol dust is presented in Rodríguez et al. (2012).

The EMEP task force of measurement and modelling (TFMM) periodically arranges intensive measurement periods (IMPs) as supplement to the continuous monitoring in EMEP (Aas et al., 2012). The third EMEP IMP took place during summer 2012 and winter 2013, and was organised in cooperation with the EU funded projects: Aerosols, Clouds, and Trace gases Research InfraStructure Network

(ACTRIS), the Chemistry–Aerosol Mediterranean Experiment (ChArME_x) and Pan-European Gas-AeroSols-climate interaction Study (PEGASOS). One of its major aims was to study the mineral dust and trace metal content of the PM₁₀ fraction.

During the third EMEP IMP, PM₁₀ filter samples were collected at 20 regional background sites (16 EMEP sites) representing different European rural background environments, by using an identical approach both with respect to sampling and subsequent speciation analysis of the mineral dust content. This qualifies for a unique data set which enables an extensive evaluation of sources, transport, and regional distribution of mineral dust across the European continent. In this paper spatial, temporal and chemical variations in mineral dust composition and trace metal concentrations are examined in order to reveal regional variations in background aerosol sources across a wide area of Europe.

2 Methods

2.1 Sampling sites

Ambient aerosol filter samples were collected at 16 EMEP and 4 other regional background sites located in 14 European countries, thus covering a wide range of the EMEP domain (Fig. 1 and Table 1). Most of these sites participated in both the summer (8 June to 12 July 2012) and the winter (11 January to 8 February 2013) IMP.

Northern Europe was represented by Aspveten (SE12) in Sweden. The Atlantic and north-western European region included Mace Head (IE31) in Ireland, Auchencorth Moss (GB48) in Scotland and Harwell (GB36) in England, all with different marine/urban air mass influences. Puy de Dome (FR30), Revin (FR09), SIRTÀ (FR20), and OPE (FR22) in France represented central western Europe; Ispra in Italy (IT04), Payerne in Switzerland (CH02), Melpitz in Germany (DE44) represented the central European area. Eastern European stations included Amberd in Armenia (AM01), Leova (MD13) in Moldova, Starina (SK06) in the Slovak Republic, and K-Pusztà in Hungary (HU02). In the south, the Spanish sites of Montsec (ES22), Montseny (ES1778), and Can Llompart, in Palma de Mallorca, (ESCL1) were taken as representative of south-western Europe, the Montelibretti (IT01) site in Italy represented central southern Europe, and Finokalia (GR02) in Crete (Greece) represented south-eastern Europe.

2.2 Sampling

Daily PM₁₀ sampling (24 h) was performed from 8 June to 12 July 2012 during the summer IMP, and from 11 January to 8 February 2013 for the winter IMP. PM₁₀ samples were collected on Teflon filters (PALLFLEX, Pall Corporation Teflon Membrane Disc Filters cod. R2PJ047, 2 μm , 47 mm, 50 pkg^{-1}) using low volume samplers (1 – 2.3 $\text{m}^3 \text{h}^{-1}$). One




Figure 1. Location of the sampling sites participating in the EMEP IMPs in summer 2012 and/or winter 2013. Blue symbols: northern Europe; purple symbols: central western Europe; black symbols: central Europe; green symbols: eastern Europe; red symbols: south-western and central southern Europe; yellow symbols: south-eastern Europe; empty symbols: mountain sites.

Table 1. Sampling sites participating in the 2012–2013 IMP. S: summer IMP; W: winter IMP. P: Pallflex.

Site	Name	Country	Altitude m a.s.l	Coordinates	Number samples	IMP	Sampler	Flow	Filter	Analysis
1	AM01	Armenia	2080	40°23′04″ N 44°15′38″ E	32	S	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
2	CH02	Switzerland	489	46°48′47″ N 05°56′41″ E	40	S 29 W	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
3	DE44	Germany	86	51°31′48″ N 12°55′48″ E	37	S 28 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
4	ES1778	Spain	740	41°46′00″ N 02°21′00″ E	41	S	Partisol/ DIGITEL	1/30 m ³ h ⁻¹	47/150 mm P	PIXE/ICPs
5	ESCL1	Spain	45	39°50′16″ N 03°01′25″ E	17	S	IND LSV	2.3 m ³ h ⁻¹	150 mm P	PIXE
6	ES22	Spain	1570	42°03′05″ N 00°43′46″ E	28	S 22 W	MCV-A/MSb	30 m ³ h ⁻¹	150 mm P	ICPs
7	FR09	France	390	49°54′28″ N 04°37′48″ E	7	S 6 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
8	FR30	France	1465	45°46′00″ N 02°57′00″ E	22	S 12 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
9	FR20	France	162	48°42′36″ N 02°08′53″ E	8	W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
10	FR22	France	392	48°33′44″ N 05°30′21″ E	7	S 9 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
11	GB36	England, UK	137	51°34′23″ N 01°19′00″ W	26	W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
12	GB48	Scotland, UK	260	55°47′36″ N 03°14′41″ W	32	S 20 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
13	GR02	Greece	250	35°19′00″ N 25°40′00″ E	39	S 30 W	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
14	HU02	Hungary	125	46°58′03″ N 19°33′11″ E	39	S 26 W	Partisol	1 m ³ h ⁻¹	47 mm P	ICPs
15	IT01	Italy	48	42°06′00″ N 12°38′00″ E	39	S 29 W	Swam 5a dual Chanel/FAI Inst.	2.3 m ³ h ⁻¹	47 mm P	PIXE
16	IT04	Italy	209	45°48′00″ N 08°38′00″ E	37	S 29 W	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
17	IE31	Ireland	5	53°19′36″ N 09°54′14″ W	31	S 14 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE
18	MD13	Moldavia	156	46°30′00″ N 28°16′00″ E	44	S 29 W	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
19	SE12	Sweden	20	58°48′00″ N 17°23′00″ E	26	S 29 W	Leckel	2.3 m ³ h ⁻¹	47 mm P	PIXE
20	SK06	Slovak Rep.	345	49°03′00″ N 22°16′00″ E	33/29	S 29 W	Partisol	1 m ³ h ⁻¹	47 mm P	PIXE

field blank was collected at each of the sampling sites for each IMP in addition to the analysis of laboratory blanks.

At Montseny (ES1778), PM₁₀ filters were sampled concurrently by a DIGITEL high volume sampler (30 m³ h⁻¹), loaded with Quartz microfiber filters (PALLFLEX, Pall Corporation QAO2500, 150 mm) to compare different methods for measuring mineral dust.

Two additional sites participated in the EMEP IMP, but PM₁₀ filters were sampled and analysed by different methods. At Montsec (ES22) PM₁₀ was collected on Quartz microfiber filters (PALLFLEX, Pall Corporation QAO2500, 150 mm) using a MCV/CAV-A MSb high volume sampler (30 m³ h⁻¹), whereas at K-Pusztá (HU02) PM₁₀ was collected on Quartz microfiber filters (PALLFLEX, Pall Corporation, 47 mm) by using low-volume samplers (1 m³ h⁻¹).

Information regarding sampling and filter analysis is summarised in Table 1.

2.3 Chemical characterisation

Most samples collected during these campaigns were analysed by PIXE (Particle-Induced X-ray Emission). PIXE is a useful technique for the analysis of refractory elements in atmospheric aerosol samples. The convenience of PIXE for atmospheric aerosol research has been evaluated by Lucarelli et al. (2011) and Maenhaut (2015). This technique allows the simultaneous determination of concentrations for elements with atomic number higher than 10 with a good sensitivity. PIXE is a non-destructive technique that does not require a specific sample preparation. The capability of analysing samples with very low concentrations without sample pre-treatment lowers the risk of contamination and analyte losses.

In the present study, samples and both laboratory and field blank filters from all sites (with the exception of K-Pusztá and Montsec) were analysed by PIXE with 3 MeV protons at the INFN LABEC (Laboratorio di Tecniche Nucleari per l'Ambiente e i Beni Culturali – Florence, Italy) with an external beam set-up (extensively described by Lucarelli et al., 2014). At LABEC, simultaneous high sensitivity detection of most mineral elements (Na, Mg, Al, Si, K, Ca, Ti, Mn, Fe, Sr, Zr) in a filter sample can be performed within a rather short time frame (30 s to 3 min), depending on the aerosol load. The minimum detection limit for each element measured by PIXE is listed in Table S1 in the Supplement.

Samples collected by high-volume samplers at ES1778, ES22 and HU02 were analysed by ICP-AES and ICP-MS at the IDAEA CSIC laboratories, following the method devised by Querol et al. (2001). At ES1778, the correlation between the concentrations obtained by PIXE and ICP-AES/MS was very high for most major and trace elements (Al, S, K, Fe, Ca, Ti and Mn), with $R^2 = 0.95\text{--}0.99$ and slopes close to 1 (1.01–1.05). Higher slopes were determined for Na (1.13) and Mg (1.26), although $R^2 > 0.9$. Correlations were lower for V, Cr, Pb and Sr ($R^2 = 0.60\text{--}0.69$; slopes 0.3 to 0.9), and very low

for Ni ($R^2 = 0.19$, slope 0.4; see Fig. S1 and Table S2 in the Supplement). For all those elements, concentrations were close to the MDL in PIXE.

The content of carbonate carbon was determined for 86 filter samples collected at 6 sites; i.e. ES1778, IT01, IT04, CH02, GR02 and MD13. The analysis followed the approach by Karanasiou et al. (2011), which by acidification (phosphoric acid) transforms the filter sample CO₃²⁻ content into CO₂. CO₂ is subsequently determined by a flame ionisation detector of a commercial thermal-optical analyser, after reduction to CH₄. The method detection limit was 0.2 μg C cm⁻².

2.4 Data treatment

Some elements such as Na, Mg, Ca and K are associated with both mineral dust and sea-salt aerosol. The sea-salt contribution was estimated for each element prior to estimating the mineral load. The concentrations of these elements in sea water are well known and therefore it is possible to estimate the marine contribution to their content in PM once we know the concentration of one of these elements. In marine-influenced and background areas, Cl⁻ has a major marine origin, therefore the marine fraction of each element could be estimated from Cl⁻ concentrations obtained for each sample. However, this method may result in an underestimation of the marine aerosol due to the interaction (in the atmosphere or on the filter) of NaCl with acidic species resulting in the formation of HCl or NH₄Cl, which may be volatilised in the atmosphere (Harrison and Pio, 1983).

It is well known that Na aerosols commonly have a marine origin, as halite associated with Cl⁻; however they can also derive from a range of other minerals such as clays, carbonates, feldspars and sulfates. Thus, assuming an exclusively or even dominantly marine origin for Na can cause significant errors, particularly in the southern European countries that are frequently affected by dust events. Average Na / Cl ratios obtained for each site during the summer EMEP IMP varied from 0.5 for Mace Head (IE31), between 1 and 2 for GB48, IT01 and GR02, and from 3 to 9 for the remaining sites. Therefore, given that the Na / Cl sea water ratio is 0.56 (Drever, 1997), there is a clear excess of Na (mineral or anthropogenic) and/or a depletion of Cl⁻ (by volatilisation) for most sites, except for IE31.

The mineral fraction of Na was estimated from the content of Al by using the ratio determined by Moreno et al. (2006) for soils and dust in northern Africa. Thus, the mineral sodium (non-sea-salt sodium, nssNa) was obtained by multiplying Al concentration by 0.12 ($\text{nssNa} = [\text{Al}] \times 0.12$). Hence, the sea-salt fraction of Na (ssNa^+) can be estimated subtracting nssNa from the total Na.

The sea-salt fraction of calcium, magnesium, potassium and sulfate was estimated by using their sea water ratios with respect to ssNa^+ (Drever, 1997; Nozaki, 1997). Finally, the total sea-salt load was determined by the sum of Cl⁻, ssNa^+ , ssCa^{2+} , ssMg^{2+} , ssK^+ and ssSO_4^{2-} . The non-sea-salt (nss)

fractions of Ca, Mg, Mn, Na, K and SO₄²⁻ were obtained by subtracting the previously calculated sea-salt fraction from their bulk concentration.

Correlation between nssK and Al concentrations for southern sites during the summer Saharan dust events (SDEs) permitted the identification of two major sources for nssK: a mineral (dustK) and a biomass burning (bbK) source. The mineral fraction is estimated from Al (dustK = 0.31 × Al; Moreno et al., 2006), and the biomass fraction by the difference: bbK = nssK – dust K.

Finally, the total mineral dust concentration was determined by addition of the concentrations of all mineral-related elements expressed as oxides, such as Al₂O₃, SiO₂, Fe₂O₃, TiO₂, P₂O₅, nssCaO, nssMgO, nssNa₂O, and dustK₂O.

Concentrations of elemental sulfur (S) were measured by PIXE. Assuming that most S is present as sulfate in PM, concentrations of S, as measured by PIXE, were converted to SO₄²⁻ multiplying by 2.995.

2.5 Saharan dust events (SDEs)

The impact of African dust outbreaks on air quality over Europe during the periods of the study was traced by using publicly available information: i.e. dust maps supplied by the Navy Aerosol Analysis and Prediction System (NAAPS) model from the Navy research Laboratory (<http://www.nrlmry.navy.mil/aerosol>; data and/or images from the BSC-DREAM8b (Dust REgional Atmospheric Model) model, operated by the Barcelona Supercomputing Center (BSC, <http://www.bsc.es/projects/earthscience/BSC-DREAM/>); and NASA Terra – MODIS satellite imagery (<http://modis.gsfc.nasa.gov/>).

During the summer 2012 IMP, two African dust outbreaks occurred: the first from the 17 to 23 June and the second one from 28 June to 7 July. The two episodes were initiated by the intense heating of the Sahara and the consequent development of the north African thermal low south of the Atlas Mountains, coupled with anticyclonic conditions at upper atmospheric levels. This is the most frequent scenario causing dust outbreaks over south-western Europe (Moulin et al., 1998; Rodríguez et al., 2001; Escudero et al., 2007; Pey et al., 2013), with the convective system pumping dust up to 5000 m a.s.l. Once the dust is injected into the mid-troposphere it may be transported toward western Europe by the eastern branch of the high (pressure) present over northern Africa (Rodríguez et al., 2001). In such cases, the air masses are heavily loaded with dust and are transported toward the north, covering most of the western Mediterranean basin, forming a wide plume of dust. As shown in the satellite image (Fig. S2b), by 25 June 2012 Saharan dust had spread to the west across the Canary Islands and Madeira. The dust started blowing a couple of days earlier in Algeria and Mali, and travelled hundreds of kilometres toward the north-west. Over the Atlantic Ocean, the dust made a giant turn toward the east, in the direction of the Mediterranean Sea. As shown

by the NAAPS model, the plume first hits the Iberian Peninsula, moving north and eastwards successively affecting parts of central, south central, and south-eastern Europe (Fig. S2a).

Short but intense dust episodes occurred during the winter 2013 IMP, affecting mainly sites in the eastern Mediterranean. This is the typical scenario in winter, when the development of low pressure systems gives rise to the rapid transport of dust at surface levels towards the east (Moulin et al., 1998), resulting in severe dust episodes in the eastern Mediterranean region (Pey et al., 2013). As shown in Fig. S3, a dust plume blew off the coast of Libya and crossed the Mediterranean towards the east on 20 January, reaching Greece and eastern Europe by 21 January. Other episodes impacted the eastern Mediterranean on 3 and 7 February. The satellite image corresponding to 7 February reflects the occurrence of multiple dust plumes transported from the coast of Libya towards the northeast (Fig. S3). The transport at surface levels results in a low spatial dispersion of the plume.

3 Results

3.1 PM₁₀ levels

Concentrations of PM₁₀, measured by gravimetry or by automatic monitors (e.g. TEOM) were available for the samples analysed at most sites, with the exception of FR20, FR30 in winter and FR22 in summer. PM₁₀ concentrations in summer were, in general, higher in the southern sites (Fig. 2). Mean levels ranged from 20 to 31 µg m⁻³ at DE44, ES22, GR02 MD13, ES1778, and IT01; from 12 to 20 µg m⁻³ at CH02, FR09, AM01, IT04, HU02, SK06 and ESCL; and from 3 to 8 µg m⁻³ at FR30, GB48 and SE12. The high levels measured at IT01 are probably due to its proximity to the city of Rome, whereas similarly high levels registered in ES1778 are attributed to the occurrence of two African dust outbreaks (see below).

In winter, low PM₁₀ levels were registered in northern and south-western Europe (3–9 µg m⁻³) and the highest PM₁₀ levels were measured in southern, eastern and central Europe (20–40 µg m⁻³). High PM₁₀ levels were recorded along a north–west to south–east transect, ranging from 22 to 40 µg m⁻³, with the highest concentrations observed for GR02, CH02 and IE31 (40, 37 and 36 µg m⁻³, respectively).

3.2 Spatial distribution and elemental affinities

Average concentrations of major and trace elements determined for each site during the summer and winter IMPs are summarised in Tables S3 and S4. In the current section, the spatial variation of major elements is discussed. The elements were classified according to their main affinities into marine, mineral, or mixed origin groups.




Figure 2. Mean PM₁₀ concentrations ($\mu\text{g m}^{-3}$) recorded during the June/July 2012 and January/February 2013 EMEP IMPs. The diameter of the circles is proportional to the concentrations.

3.2.1 Sea-salt-related elements

Sea salt (estimated as described in the methodological section, Sect. 2.4) accounts for a larger fraction of PM₁₀ at coastal sites, around 10% on average, with important variations in winter (Fig. 3). During the summer IMP, the average sea-salt load ranged from 1.9 to 2.7 $\mu\text{g m}^{-3}$ at the coastal sites (10% of PM₁₀ at ESCL1 and GR02 in the Mediterranean, 16% at IE31 in the Atlantic), from 0.2 to 0.7 $\mu\text{g m}^{-3}$ at sites situated close to the coast (4–10% of PM₁₀ at IT01, GB48, ES1778 and SE12), and were below 0.2 $\mu\text{g m}^{-3}$ for the remaining sites. In winter, the sea-salt load was up to 1 order of magnitude higher, compared to the summer ones at the north-western sites, reaching an average concentration of 19 $\mu\text{g m}^{-3}$ at the coastal IE31 (>50% of PM₁₀) and values greater than 2 $\mu\text{g m}^{-3}$ at GB48 and GB36 (30 and 14% of PM₁₀, respectively). In the Mediterranean area, the average sea-salt contribution was slightly higher at GR02 (3.9 $\mu\text{g m}^{-3}$, 10% of PM₁₀) compared to the other sites (<1 $\mu\text{g m}^{-3}$).

3.2.2 Mineral dust

In order to identify major sources of mineral dust in the study area, we have explored the correlations between elements with a major mineral affinity. We find that a number of mineral-related elements correlate strongly with Al and Si, whatever the site, suggesting a dominant aluminium-silicate (Al-Si) occurrence in PM. These components are Si, Ti, and Fe, and all showed higher concentrations at the southern and eastern sites (as deduced from data shown in Tables S3 and S4).

During the summer IMP, concentrations of Al were markedly higher in the south-western, south central, and eastern European regions, reaching average concentrations above

1 $\mu\text{g m}^{-3}$. These decreased northwards, being <0.1 $\mu\text{g m}^{-3}$ at the north-western and northern European sites. High levels of Al recorded at the southern sites were related to the impact of Saharan dust events (as will be discussed in the following sections) and to resuspension of soil dust on a regional scale. In eastern Europe, where impact of Saharan dust events in PM levels was not as clear, a high contribution of regional/local dust is deduced. Concentrations of Al decreased at most sites in winter, being <0.2 $\mu\text{g m}^{-3}$ in the south and <0.1 $\mu\text{g m}^{-3}$ in the north. An exception was seen for GR02 (average of 2 $\mu\text{g m}^{-3}$) due to the impact of Saharan dust events even in winter. A similar spatial distribution was observed for Si, given its very high correlation with Al ($R^2 = 0.98$).

Figure 4 depicts cross-correlation plots between the R^2 coefficients and the slopes of the regression equation ($y = ax + b$) calculated for major elements at each site. The figure shows that there was a high correlation ($R^2 > 0.8$) between Si and Al (excluding IE31, $R^2 = 0.23$) when considering both IMPs, with $R^2 > 0.95$ at most sites. The slope of the equation ($\text{Si} = x\text{Al} + b$) was within 1.9 (IT04 and DE44) and 2.9 (MD13), ranging between 2.0 and 2.6 for most sites. Thus, the average Si / Al ratio was 2.34. There was a spatial gradient, this being most evident in summer, which showed a downward trend of the Si / Al ratios from eastern (slope = 2.4–2.9) towards western Europe (2.0–2.2). These high correlation coefficients and the Si / Al ratios point to a clay dominated source (mainly illite, Si / Al = 3; Scheuven et al., 2011) for both elements. The relatively low Si / Al ratio seen for southern and south-western Europe was probably due to a higher contribution of kaolinite (Si / Al = 1.0). In winter, this spatial variability is not as clear, possibly due to the lower concentrations.




Figure 3. Spatial distribution of the mean sea-salt aerosol concentration ($\mu\text{g m}^{-3}$) and its relative contribution (%) to PM₁₀ during the EMEP IMP in summer 2012 and winter 2013. The diameter of the circles is proportional to the concentrations and percentages.

Aluminium also correlates well with Fe, with $R^2 > 0.70$ for most sites, pointing to a major aluminium-silicate association. The determination coefficient is $R^2 > 0.90$ for some sites when considering both IMPs (AM01, GR02, CH02, MD13, SK06, ES22, ES1778, IT01, FR30, FR22), confirming a mostly unique clay-related source for Fe at these sites. In these cases the slope varies from 0.5 (ES22, ES1778, FR30, FR22, SK06) to 0.7 (AM01, MD13) reflecting a different local mineral assemblage. Lower correlations were obtained for IE31 ($R^2 = 0.01$) and IT04 ($R^2 = 0.15$). At the other sites, R^2 ranged from 0.6 to 0.8; hence a significant contribution of other mineral or non-mineral sources can be suspected. The correlation was slightly higher in summer, with R^2 ranging from 0.7 to 1.0 at all sites, except at IE31 ($R^2 = 0.10$), reflecting the higher contribution of mineral dust. In winter, however, R^2 is lower than 0.8 for most sites, with slopes frequently > 1 , reflecting a lower contribution of mineral dust and probably a higher contribution of anthropogenic sources of Fe.

Calcium has a major mineral affinity, and is only a minor species in sea salt, being present in the atmosphere as carbonate (calcite CaCO_3 , dolomite $\text{CaMg}(\text{CO}_3)_2$) or as calcium oxide (CaO), and to a minor extent as calcium sulfate (gypsum – $\text{CaSO}_4 \times 2\text{H}_2\text{O}$, and/or anhydrite – CaSO_4) or as silico-aluminates (Ca-plagioclase) (Scheuven et al., 2013). This element is usually related to natural sources (soil resuspension), although it can be emitted by a number of anthropogenic sources, such as road dust and construction activities (Amato et al., 2009). Calcium carbonate may interact with acidic compounds in the atmosphere forming coarse secondary calcium nitrates ($\text{Ca}(\text{NO}_3)_2$) and calcium sulfates ($\text{CaSO}_4 \times x\text{H}_2\text{O}$) (Dentener et al., 1996; Krueger et al., 2004; Alastuey et al., 2005; Hwang and Ro, 2006). The samples analysed for carbonate Carbon (CC) had a rather consistent CC/Ca ratio considering all sites and regardless of the samples being affected by African dust episodes or not (Fig. 5). When including all samples, the overall CC/Ca ratio was 0.12 ± 0.01 ($R^2 = 0.75$) and the




Figure 4. Scatter plots of determination coefficient vs. slope for concentrations of selected elements at the sampling sites.




Figure 5. Cross correlation plot between carbonate carbon (CC) and nssCa content (in $\mu\text{g m}^{-3}$) in PM₁₀ at sites in different regions.

intercept $-0.03 \pm 0.01 \mu\text{g CC m}^{-3}$. Assuming a method recovery of 100% (i.e. all CC present on the filter sample is determined), the CC/Ca ratio value suggests that at least half of the Ca is present in PM₁₀ in other forms than mineral car-

bonate, e.g. $\text{Ca}(\text{NO}_3)_2$ and $\text{CaSO}_4 \times x\text{H}_2\text{O}$. Given the high correlation between Ca and CC, it can be deduced that non-carbonate Ca follows from the reaction between carbonates and acidic compounds.

3.2.3 Elements with a mixed source origin

Magnesium has a mixed origin and is associated with both marine aerosol and mineral dust. A dominantly marine origin was deduced for Mg at the northern sites (IE31, SE12 and GB48), this element being highly correlated with Na ($R^2 = 0.81\text{--}0.98$) and with a Na/Mg ratio ranging from 5.2 to 6.7, slightly lower than the ratio for sea salt (8.4; Drever, 1997). A mixed marine/crustal origin was deduced for southern and central Europe with two different correlations between Na and Mg for days with dust or marine influence. The nssMg may be associated with clays, carbonates (dolomite), aluminium-silicates or salts (Scheuvs et al., 2011). Considering all samples, the concentrations of nssMg were highly correlated with Al ($\text{nssMgO} = 0.30 \times \text{Al}_2\text{O}_3$, $R^2 = 0.76$). However, the correlation between nssMg and Al varied considerably between sites, reflecting different mineral associations for nssMg. Three groups of sites can be distinguished as a function of R^2 . The determination coefficient was <0.3 at IE31, GB48 and SE12; >0.9 at AM01,

CH02, GR02, MD13, SK06, ES1778 and FR30; and between 0.6 and 0.9 at the remaining sites. For most sites, the slope ranged between 0.15 and 0.2 (Fig. 4). At two eastern sites AM01 and GR02, a unique crustal origin was deduced for nssMg, and with relatively high concentrations. In this region the Mg / Al ratio increased to 0.5 ($R^2 = 0.98$), which could be related to the high Mg geochemical anomaly of the region. An intermediate slope (0.25) was obtained for MD13.

Non-sea-salt potassium (nssK) has a major aluminium-silicate affinity and may be present in minerals such as K-feldspars, muscovite and illite (Scheuvens et al., 2013), but can also be emitted during biomass combustion (Andreae, 1983; McMeeking et al., 2009). When considering all samples, nssK and Al concentrations were moderately correlated ($R^2 = 0.47$), with a slope of 0.26 and a low intercept value ($0.1 \mu\text{g m}^{-3}$). In summer, the correlation was higher ($R^2 = 0.78$), indicating a major Al-Si affinity for nssK. Figure 4 shows this correlation varies considerably from site to site, especially in winter. Most southern sites (ES1778, GR02, ESCL1, ES22, AM01, FR30) were characterised by high R^2 (0.8–1.0) and slopes between 0.2 and 0.3, confirming a major mineral origin. For northern and central European sites, as well as for some southern sites (IT01), the determination coefficients were lower ($R^2 < 0.6$). In summer, this grouping is obvious, but the slopes ranged 0.15–0.4 and $R^2 = 0.5$ –1 (except for IE31) for all sites. In winter, there is a higher dispersion (Fig. 4). High nssK / Al correlation ($R^2 = 0.8$ –1) was only determined for ES1778, FR30 and GR02, with slopes 0.3–0.5. At the remaining sites $R^2 < 0.6$ (ranging 0–0.6) and the slopes varied from 0.1 to 7. This indicates an additional potassium source in winter, which most likely is biomass combustion. This additional source was more evident at certain sites (e.g. IT04, CH02, GB36, FR20, FR22, SE12; Fig. 4) where nssK / Al slopes > 1 were recorded in winter, revealing a significant contribution from biomass burning. Thus, correlation of nssK with Al permitted us to identify two major sources for nssK, namely those of dust (dustK) and biomass burning (bbK). For those sites with a high correlation ($R^2 > 0.9$) between nssK and Al, we consider that nssK is mostly associated with aluminium silicates. Given the low nssK-Al, the presence of illite or muscovite is likely (Scheuvens et al., 2011). In these cases we used the average ratio K / Al in summer to calculate dustK. This ratio varied from 0.2 at ES22, 0.3 at ES1778 and FR30, 0.4 at AM01 and FR22, and 0.5 at GR02, MD13 and SK06. For the remaining sites, we used an intermediate value (nssK / Al = 0.3), which coincides with the average ratio obtained by Moreno et al. (2006) for Saharan dust samples and with the one obtained for samples collected during SDEs in this study (see next section). Once the dustK was estimated, the biomass fraction was calculated as the difference: $\text{bbK} = \text{nssK} - \text{dustK}$.

The spatial variation of the estimated dustK and bbK average concentrations in PM₁₀ is presented in Fig. 6. Concentrations of dustK were higher in summer, ranging from

10 ng m^{-3} at the northern sites to 100 – 200 ng m^{-3} at the southern sites and $> 400 \text{ ng m}^{-3}$ at MD13. In winter, dustK ranged 5 – 50 ng m^{-3} , except at GR02 with an average concentration of 400 ng m^{-3} , which is higher than in summer (200 ng m^{-3}) due to the occurrence of Saharan dust events in winter. By contrast, concentrations of bbK were higher in winter, ranging from 50 to 500 ng m^{-3} , especially in central and eastern Europe. If we compare the relative contribution of bbK and dustK to the nssK, the dust contribution was most important in summer for the Mediterranean and the eastern European sites, whereas for the northern European sites dust and biomass were equally large sources. In winter, bbK clearly dominated, reflecting the impact of biomass combustion mainly at the northern and central sites. However, this estimation may be subject to significant errors for specific cases. Thus, the relatively high concentrations of bbK estimated for IT01 and MD13 in summer could be due to a different local soil composition.

These findings demonstrate the influence of local (both natural and anthropogenic) and external emissions as revealed by the ratios between PM mineral components. These influences can result in significant errors when trying to estimate the PM mineral load by applying factors to the concentration of a single measured element, as proposed by a number of previously published works, an approach that should therefore be used with caution.

3.3 Mineral dust contribution: impact of Saharan dust events

Figure 7 shows the spatial variation of the average mineral dust load contribution determined for each site and IMP following the procedure explained in Sect. 2.4. In general, higher concentrations of mineral dust were recorded in summer compared to winter. The exceptions were GR02, strongly affected by Saharan dust events in winter, and IE31, with very low concentrations of mineral dust. The summer maxima of mineral dust were more evident in the southern and eastern countries, showing a spatial distribution similar to that described for PM₁₀ (Fig. 2). The highest dust load (5 to $10 \mu\text{g m}^{-3}$) was determined at the MD13, IT01, ES22, AM01 and ES1778 sites. Intermediate levels (between 2.5 and $5 \mu\text{g m}^{-3}$) were obtained at GR02, ESCL1, HU02, SK06, FR09, FR22 and IT04, followed by CH02 and DE44 (1.5 – $2 \mu\text{g m}^{-3}$). At the remaining sites (FR30, SE12, GB48 and IE31) the dust load was $< 0.5 \mu\text{g m}^{-3}$. Hence, the mineral load accounted for less than 10 % of PM₁₀ in summer at the northern sites (2 % in IE31, 6 % in SE12 and 9 % in DE44), for 15 to 25 % of PM₁₀ at most sites, and for more than 30 % at ES22 (34 %), MD13 (38 %) and AM01 (42 %).

The time series of the ambient air mineral dust concentration during the summer IMP is presented in Fig. 8. Mineral dust concentrations increased at southern and at certain central European sites during the two SDEs observed during the summertime IMP; i.e. from 17 to 23 June and from 28 June




Figure 6. Spatial distribution of the estimated average concentrations (in ng m^{-3}) of the mineral (dustK) and the biomass burning (bbK) potassium at each site in the summer 2012 and winter 2013 IMPs. The diameter of the circles is proportional to the concentrations.

to 7 July. The mineral dust concentration first increased at the south-western sites (ES22 and ES1778), reaching $12 \mu\text{g m}^{-3}$ on 17 June (Fig. 8). As the plume moved eastward (see Fig. S2), levels increased at the two Italian sites, reaching $15 \mu\text{g m}^{-3}$ at IT01 on 22 June. On 28 June a second and more intense plume was observed over the European continent, spreading to distant areas such as Germany and the British Isles. The second SDE first impacted the Iberian Peninsula, which experienced daily mineral dust concentrations of $34 \mu\text{g m}^{-3}$ at ES22 and $20 \mu\text{g m}^{-3}$ at ES1778 on 28 June. Mineral dust levels decreased northwards, reaching maximum daily concentrations on 28 June at CH02 ($14 \mu\text{g m}^{-3}$), ESCL1 ($11 \mu\text{g m}^{-3}$), FR09, FR22 and DE44 ($7\text{--}8 \mu\text{g m}^{-3}$), FR30 and GB48 ($3 \mu\text{g m}^{-3}$). On 29 June, the concentrations of mineral dust increased at ES22 (up to $39 \mu\text{g m}^{-3}$) and at ES1778 and started increasing at the Italian sites (IT01 and IT04). On this day, the concentration of mineral dust decreased at the mentioned central and north European sites (CH02, ESCL1, DE44, FR30 and GB48). On 30 June, the mineral load started decreasing at ES22 but still increased

at ES1778, reaching $35 \mu\text{g m}^{-3}$. Subsequently, the mineral load decreased at these sites, but increased at IT04 and IT01, reaching $9 \mu\text{g m}^{-3}$ on 30 June, reflecting the transport of the dust plume towards the eastern Mediterranean region. The first of July, levels of mineral dust reached the maximum at ES1778, IT01 and HU02 ($12\text{--}14 \mu\text{g m}^{-3}$). On 5 July the mineral load peaked at GR02, reaching $15 \mu\text{g m}^{-3}$. From 1 July and until the end of the month in 2012 EMEP IMP, levels remained relatively high in the central and eastern parts of the Mediterranean Basin. During these dust episodes, the mineral dust concentration was always higher at ES22 than at ES1778, reflecting the transport of dust at high altitudes in summer. Finally, during the SDEs, the mineral dust contribution to PM₁₀ was on average 35 % at the affected sites, ranging from 25 % at DE44 to 55 % at ES22.

It should be mentioned that at the Spanish sites the contribution of the Saharan mineral dust during the sampling period accounts for two exceedances of the daily limit value of $50 \mu\text{g PM}_{10} \text{m}^{-3}$, established by the EC Directive (EC, 2008).




Figure 7. Spatial distribution of the average mineral dust concentration ($\mu\text{g m}^{-3}$) and its relative contribution (%) to PM₁₀ obtained at each site in the summer 2012 and the winter 2013 IMPs. The diameter of the circles is proportional to the concentrations and percentages.

The SDEs only partly affected eastern Europe. Thus, the mineral dust loading observed in this region is presumably related more to the influence of local and/or regional sources. The presence of distinctive geochemical ratios (e.g. Si / Al and Mg / Al) characteristic for this region is confirmed in the present study which has allowed us to demonstrate the importance of crustal sources in the eastern European region. Further research is needed to identify these sources in more detail, e.g. to establish emission factors, and to quantify their impact on ambient PM levels over longer periods. Another complicating factor is that this area is likely impacted at times by emissions of dust from the Arabian deserts.

In winter, the concentrations of mineral dust were lower for all sites except for GR02 (Fig. 7), where an average concentration of $13.5 \mu\text{g m}^{-3}$ was calculated due to the influence of short but intense Saharan dust events (see Fig. S3). Levels between 1 and $2 \mu\text{g m}^{-3}$ were measured at IT04, IT01, MD01, and FR20. At the remaining sites, the dust load was $< 1 \mu\text{g m}^{-3}$. In contrast to the summer IMP, the dust load did not reflect the temporal variation of PM₁₀ during the winter

IMP. For central and northern Europe, the mineral fraction accounted for less than 5 % of PM₁₀ at most sites, whereas it ranged between 5 and 10 % for the southern and eastern sites. The 34 % contribution of mineral dust to PM₁₀ at GR02 was an exception. The lower mineral dust loads in winter were attributed to a lower impact of the Saharan dust outbreaks (affecting only the eastern part of the Mediterranean) and reduced soil resuspension in this period.

Figure 9 shows the time evolution of the mineral dust concentration calculated for PM₁₀ during the winter IMP. The dust load was low at most sites, but peaked frequently at GR02 due to the impact of short but severe dust episodes which severely deteriorate air quality in the eastern Mediterranean area (Dayan et al., 2008; Querol et al., 2009; Pey et al., 2013). Thus, daily concentrations of mineral dust $> 50 \mu\text{g m}^{-3}$ were recorded at GR02 over 3 days (ranging $60\text{--}83 \mu\text{g m}^{-3}$). The lower spatial dispersion of most winter episodes is a consequence of the transport at surface levels. During 21 January a dust plume blew off the coast of Libya and crossed the Mediterranean towards the east (see




Figure 8. Time series of mineral dust concentrations during the June/July 2012 EMEP IMP at sites affected by Saharan dust outbreaks. Two major dust episodes were observed in June; in both cases, the dust plume moved along a west to east transect.




Figure 9. Time series of mineral dust concentrations during the January/February 2013 EMEP IMP at selected sites affected by the Saharan dust outbreaks. Levels of mineral dust at GR02 (right axis) reflect the impact of short but intense dust pulses.

Fig. S3). This plume extended to areas in northern Italy and dust concentrations increased simultaneously at distant sites such as GR02, HU02, SK06 and DE44, although it is difficult to attribute this increase exclusively to the impact of the dust plume.

3.4 Local/regional mineral dust contribution: chemical composition

In the current section, days not impacted by Saharan dust were studied to characterise the chemical profiles of local dust sources and to quantify their contribution to PM. Average dust concentrations were calculated for the two measurement periods without impact of Saharan dust (NO-SDE, Fig. S4). The results show a clear spatial pattern, as the dust concentration decreases towards the north and the west. Hence, dust attributed to regional/local sources accounted for more than 25 % of PM₁₀ at the eastern sites MD13 (7 µg m⁻³) and AM01 (6 µg m⁻³; summer only). At the southern sites (western, central and eastern Mediter-

anean) local/regional dust ranged between 1.5 and 4 µg m⁻³, accounting for 8–15 % of PM₁₀. The higher contribution at IT01 (4 µg m⁻³, 15 % of PM₁₀) is likely related to anthropogenic sources in the urbanised areas nearby the site. At most central European sites, the mineral dust concentration ranged between 0.5 and 2 µg m⁻³, accounting for 5–10 % of PM₁₀. Lower values were estimated for sites in northern and western Europe.

The chemical composition of mineral dust may vary considerably depending on the location, soil composition and influence of external sources (i.e. Saharan dust). As shown in Sect. 3.3, Saharan dust had a significant influence on PM₁₀ levels at the southern sites during the summer IMP. As dust can be mixed with other sources and transformed during transport, the current set of data provides a unique opportunity to investigate the impact of Saharan dust on the chemical composition of mineral dust and on the variation of dust composition during transport.

Chemical composition of the mineral dust, expressed as oxides, was averaged at each site for those days with a significant impact of Saharan dust (Fig. 10). During the winter IMP only GR02 was considered, as it was the only site clearly impacted for a significant period (more than 1 day), although other sites such as HU02, SK06, and MD13 were also briefly impacted. Figure 10 shows a steep gradient in the concentration of mineral dust dominated by African dust, with the highest values seen for the areas located close to the source region; i.e. the eastern Mediterranean (in winter) and the southwest of Europe (summer). The relative composition (expressed as %, Fig. 10 bottom) is quite similar among all sites, with differences for GR02 in winter (GR02-W), IT01 and ESCLI, which have a larger contribution of CaO.

Ternary diagrams with average dust composition during Saharan episodes are presented in Fig. S5. With respect to SiO₂, Al₂O₃ and Fe₂O₃ (Fig. S5), the average composition was almost identical for all sites, indicating a similar silico-aluminous composition. However, for similar SiO₂/Al₂O₃ ratios, the contribution of CaO varies (Fig. S5), being higher during the winter (GR02-W) events. This is probably related to the different source areas for the summer and winter periods. In summer (see Figs. S2 and S3), dust was emitted from southern Algeria and Mali, whereas in winter dust blew off the coast of Libya. This is in agreement with previous studies showing higher content of Ca-carbonates for dust coming from north-eastern Africa (Formenti et al., 2011; Scheuven et al., 2013). Relatively higher contents of CaO were also measured at ESCLI and IT01. This could be related to local contribution from soil resuspension given that the geology in these areas is characterised by the presence of carbonate rocks (limestone). At IT01, an important fraction of calcium could be also related to the contribution of local anthropogenic sources, given its proximity to Rome.

During the summer episodes, southern sites (orange and red colours) tend to have a higher content of CaO and MgO and a relatively lower content of K₂O and Fe₂O₃ compared to




Figure 10. Average concentration of mineral dust components (absolute values – $\mu\text{g m}^{-3}$, top) and the relative contribution of components (% mineral load, bottom) in mineral dust at selected sites when affected by African dust episodes. Results from both summer and winter (GR02-W) IMPs are included.

central and northern European sites. This could be attributed to the different composition of local dust or to the preferential settling during transport due to the particle size and morphology of specific minerals. Again, higher contents of CaO were obtained for IT01, GR02-W and ESCLI whereas ratios of MgO/CaO keep constant for GR02-W and ESCLI, indicating a soil-related source (carbonates). CaO/MgO was higher at IT01, probably due to an anthropogenic contribution (construction, demolition, road dust).

Ternary diagrams based on the average composition of samples not affected by Saharan dust are shown in Fig. S6. As for the African dust episodes, the $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio was fairly constant for all sites except IE31. The possibility of the low levels observed at IE31 affecting the results should not be excluded. For the other components, there is a wider variation in comparison with the African episodes (see Fig. S5). When plotting SiO_2 , Al_2O_3 and CaO, the contribution of CaO clearly decreases from the southern to the eastern and northern sites. The obvious presence of this trend in the high altitude sites (ES22, AM01 and FR30) emphasises that it is a likely marker for far-travelled particles derived from the local/regional geology in the source areas. An inverse pattern is observed when plotting SiO_2 and Al_2O_3 with either Fe_2O_3 or K_2O ; there is a clear increasing trend of the relative contribution of K_2O and Fe_2O_3 from the southern to the eastern and northern sites. However, in this case the average concentrations are very similar at the three high altitude sites, suggesting that this trend reflects the different influence of other sources such as biomass burning for K, or iron steel industry for Fe_2O_3 .

3.5 Sulfate

Sulfur is typically present in ambient air as sulfate (SO_4^{2-}). Therefore, although PIXE measures concentrations for S, we have chosen to use sulfate in the present text. Non-sea-salt sulfate is a major secondary component formed by the oxidation of SO_2 , mainly emitted by anthropogenic sources such as fossil-fuel combustion processes and metal smelters, as well as natural sources. Moreover, non-sea-salt sulfate (nssSO_4^{2-}) may have a minor mineral association, mostly as coarse gypsum, and can also be released from marine biogenic and volcanic emissions (Bates et al., 1992 and references therein).

Concentrations of nssSO_4^{2-} differed both spatially and temporally (Fig. 11). The highest mean SO_4^{2-} concentration in summer was observed at the south-eastern site GR02 ($5.8 \mu\text{g m}^{-3}$). Relatively high concentrations were also determined for the eastern, south central, and south-western sites, ranging between 2.6 and $3.4 \mu\text{g m}^{-3}$. Lower levels were measured at central and northern sites ($0.5\text{--}1.1 \mu\text{g m}^{-3}$) with an intermediate concentration for DE44 ($2.0 \mu\text{g m}^{-3}$). In winter, higher concentrations of SO_4^{2-} were observed in eastern and central European sites (ranging $2\text{--}3.8 \mu\text{g m}^{-3}$), whereas the average concentration was $<2 \mu\text{g m}^{-3}$ for southern and northern Europe.

Given the significant impact of Saharan dust outbreaks in southern European countries, it can be speculated that a minor fraction of SO_4^{2-} could be primary gypsum, despite that it is a minor (usually $<2\%$) component of Saharan dust with particularly low concentrations in most north African source areas (Claquin et al., 1999; Scheuvens et al., 2013; Journet et al., 2014). To identify the contribution of primary gypsum to the observed concentrations of SO_4^{2-} , we have investigated the correlation of SO_4^{2-} with major mineral elements. Figure 12 presents ternary diagrams for average concentrations of SiO_2 , Al_2O_3 and SO_4^{2-} for SDE and NO-SDE samples. For SDE the spatial pattern was different from that observed for Ca (Fig. S5). The highest relative SO_4^{2-} contributions were obtained for GR02-S (summer) and central European sites, with lower contributions for GR02-W (winter) and some southern European sites, whereas higher relative contributions of Ca were determined for GR02-S and southern sites. A similar but opposite spatial trend of Ca was observed during NO-SDE (Fig. 12, right, and Fig. S6). The low concentrations of SO_4^{2-} at the mountain sites, indicated a low contribution of primary gypsum. The increase in concentrations of SO_4^{2-} at distant sites from northern Africa even during SDE suggests a major anthropogenic source, most probably related to the impact of regional SO_2 emissions, and originating mainly from fuel oil and coal combustion. A major source of SO_4^{2-} in the Mediterranean region can be related to shipping (heavy oil) emissions, whereas in central and eastern Europe high concentrations are more often related to stationary sources, such as individual coal heated




Figure 11. Spatial distribution of the mean nss-sulfate concentrations ($\mu\text{g m}^{-3}$) during the summer 2012 and winter 2013 EMEP IMPs. The diameter of the circles is proportional to the concentrations.




Figure 12. Ternary diagram for major mineral dust components (SiO_2 and Al_2O_3) and SO_4^{2-} for days with (left, SDE) and without impact of SDE (right, NO-SDE). Orange: GR02; red: south-western and central southern Europe sites; purple: central western Europe; black: central Europe sites; blue; northern Europe and Atlantic sites; green: eastern Europe sites; empty symbols: high altitude sites.

ovens and coal power plants, which are discussed in more detail in the next section.

3.6 Trace elements

Figures 13 and 14 show the spatial variation of selected trace elements during the summer and winter IMPs. The concentrations varied substantially between sites, whereas there was typically a minor difference with respect to concentration between summer and winter at most sites.

3.6.1 Mineral-related elements

Some metals, such as Ti (shown in Fig. 13), Sr and Rb, were highly correlated with major mineral elements owing to their dominantly crustal origin. Consequently, these elements showed a spatial distribution similar to that of mineral dust, with higher mean concentrations in the southern and eastern

European sites during summer, and at GR02 during winter. Mean concentrations of these crustal-related elements were typically substantially lower in winter compared to summer. High concentrations of Sr at IE31 in winter were related to its partial marine origin, whereas its presence at IT04 was probably related to resuspension of road dust. The highest Sr concentrations in summer were observed at IT01, reflecting the influence of the anthropogenic emissions from the nearby city of Rome. In general, having a major carbonate affinity, Sr concentrations reflect the spatial distribution of Ca.

3.6.2 Combustion-related elements

Vanadium (V) is a typical tracer of heavy oil combustion, usually associated with nickel (Ni) and SO_4^{2-} (Viana et al., 2008; Alleman et al., 2010). Elevated concentrations ($5\text{--}6\text{ ng m}^{-3}$) of V were recorded at the southern sites in




Figure 13. Spatial distributions of the mean concentrations (in ng m^{-3}) of Ti, V, and Ni, determined at each site during the summer and the winter EMEP IMPs. The diameter of the circles is proportional to the concentrations.

summer, whereas V was $<2 \text{ ng m}^{-3}$ at the remaining sites. Concentrations of V decreased in winter, being relatively higher at coastal sites such as IE31 (1.9 ng m^{-3}) and GR02 (3.4 ng m^{-3}). For the remaining sites, the mean V concentration was between 0.5 and 1 ng m^{-3} . Nickel is commonly associated with V and is used to trace combustion of fuel oil; however it is also emitted by metallurgical processes (iron and steel manufacturing) (Viana et al., 2008; Pandolfi et al.,

2011). In summer, the highest levels of Ni were recorded at southern (2 – 2.5 ng m^{-3}) and eastern European sites (1 – 2 ng m^{-3}), with a maximum value of 3.3 ng m^{-3} at HU02, being $<1 \text{ ng m}^{-3}$ at the other sites. In winter, average concentrations were lower, with maximum values at the Italian sites (1.2 – 1.9 ng m^{-3}). At the other sites, the concentration ranged between 0.4 and 1.1 ng m^{-3} , whereas it was $<0.4 \text{ ng m}^{-3}$ at the mountain sites.




Figure 14. Spatial distributions of the mean concentrations (in ng m^{-3}) of Cr, As, and Cu, determined at each site during the summer and the winter EMEP IMPs. The diameter of the circles is proportional to the concentrations.

Figure S7 shows the correlation between the average concentrations of V, Ni, and SO_4^{2-} , the V/Ni ratio obtained for the two IMPs, and the correlation coefficients and the slopes of the regression equation calculated for these elements at each site. At the Mediterranean sites (red and orange colour), which experienced the highest concentrations of V, correlations were significant ($R^2 > 0.4$, $p < 0.001$) when consider-

ing V and SO_4^{2-} , as was V vs. Ni ($R^2 > 0.4$, $p < 0.0001$). The V/Ni ratio ranged from 2.3 to 2.5 (except for IT01; 1.5), which is within the range (2–4) identified for shipping emissions reported by Viana et al. (2008), Alleman et al. (2010) and Pandolfi et al. (2011). At AM01, V was found to correlate with Ni and SO_4^{2-} ; however, the V/Ni and V/ SO_4^{2-} ra-

tios were totally different from those obtained at the Mediterranean sites pointing to a different fuel combustion source.

Hence, a major common source of SO₄²⁻, V and Ni, likely related to fuel oil combustion emissions in the Mediterranean has been identified. As shown in Figs. 11 and 13, there is a clear seasonal trend for these elements at the Mediterranean sites, with higher concentrations in summer. This may be attributed to lower maritime traffic in winter (cruises) and the higher photochemical oxidation of SO₂ in summer. Some central and eastern European sites, showed relatively high SO₄²⁻ concentrations, but low concentrations of V, as well as a low correlation ($R^2 < 0.4$) between the two, indicating another major source for SO₄²⁻. As shown in Fig. 14, summer average concentrations of As were higher at south-eastern and central European sites (0.6–0.7 ng m⁻³) with the most elevated levels at AM01 (1.6 ng m⁻³). Lower values were recorded for western European sites, for which average concentrations ranged from 0.1 to 0.3 ng m⁻³. In winter (Fig. 13), the higher concentrations were obtained at central and eastern European sites (0.6–1.3 ng m⁻³). When considering the two campaigns (Fig. S8), higher correlation coefficients between As and SO₄²⁻ ($R^2 = 0.42$, $p < 0.0001$) were determined for DE44, with relatively high concentrations of As, and for ES22, with very low As concentrations. At the remaining sites, R^2 coefficients range 0.1–0.3. For similar R^2 , higher As concentrations were recorded at the central and eastern sites (Fig. S8). A similar trend was also observed for other pollutants such as Pb (see Tables S3 and S4). Sources of As, Pb, and SO₂ (and other volatile pollutants including Cd, Se, and Hg) are related to high-temperature processes such as coal combustion, roasting and smelting of ores in non-ferrous metal smelters (Pb and Cu-Ni production) and melting operations in ferrous foundries, among others (Pacyna, 1986; Nriagu and Pacyna, 1988). It can be concluded that there is a higher contribution of coal combustion sources in central and eastern Europe (mainly in winter), that the high levels of As registered at AM01 could be related to other sources, and that the sporadic impact of coal power plant emissions at ES22 cannot be discarded.

3.6.3 Other industrial sources and road traffic

Tracers of industrial activities, such as Cr, Mn and Ni, do not show a clear spatial distribution pattern (see Figs. 13 and 14). These elements may also have a mineral association and, as discussed for Ni, can be emitted by different processes. The higher concentrations of Cr, usually considered as a tracer for metallurgical activities (Querol et al., 2007), were measured in SK06 (4.4 ng m⁻³) in summer and at IT01 (3.7 ng m⁻³) in winter. At SK06 and AM01, concentrations show a very high correlation with Ni ($R^2 = 0.94$) confirming a common origin related to metallurgical activities. Relatively high determination coefficients between Ni and Cr were also obtained at MD13, IT04, CH02, IE31, and ES22. At the MD13 and

AM01 sites, high correlations of Cr and Ni with Mn and Cu, reinforce the link with metallurgical activities.

Although emitted in large proportion by industrial sources, Cu is often a tracer of non-exhaust vehicle emissions (Schauer et al., 2006; Amato et al., 2009). In summer this element was found to be present in higher concentrations at the southern and central European sites with average concentrations of 11 ng m⁻³ at IT01 and 8.8 ng m⁻³ at IT04 (Fig. 14). At the other sites, mean Cu concentrations mostly ranged from 1 to 4 ng m⁻³. In winter, high concentrations were again measured at IT04 and IT01 (15.9 and 5.7 ng m⁻³, respectively) and at FR20 (9.4 ng m⁻³). The lowest concentrations were measured at the mountain sites (ES22 and FR30, < 0.5 ng m⁻³).

To illustrate the differences in metalliferous tracers of emissions from major sources such as road traffic, fossil fuel combustion and industry, two ternary diagrams are proposed to compare the sites (Fig. S9). In these diagrams Cu is considered to be a tracer for traffic, V (and partially Ni) as tracers for shipping emissions, As for coal combustion emissions and Cr and Ni as industrial (metallurgy) tracers. The relative proportions of these four possible sources allow us to differentiate (1) Mediterranean sites together with IE31, with a higher influence of fuel combustion (shipping) emissions; (2) sites with a relatively higher influence of traffic (IT04, IT01 and FR20) located close to important cities or busy roads; (3) the eastern (AM01, SK06, HU02, MD13) and central (DE44) sites with a higher impact of coal combustion emissions; (4) some central and eastern sites more influenced by metallurgical activities. The mountain sites form a separate sub-group, having low concentrations of all of these tracers.

4 Conclusions

The third EMEP intensive monitoring period, conducted in summer 2012 and winter 2013, addressed the chemical speciation in PM₁₀ with a particular emphasis on mineral dust and trace metals. For the first time, mineral dust was determined in filter samples (PM₁₀) collected concurrently at a substantial number (20) of regional background sites across Europe, using a similar methodology at 18 of the sites; i.e. particle-induced X-ray emission (PIXE), conducted at the INFN LABEC of Florence, Italy. PIXE analysis allowed for the simultaneous detection of most mineral elements (i.e., Na, Mg, Al, Si, K, Ca, Ti, Mn, Fe, Sr, Zr) with high sensitivity.

The PM₁₀ mineral dust composition across Europe demonstrated the influence of both local (natural and anthropogenic) and external sources, as evidenced by ratios of different mineral components in the PM filter samples. In general, higher concentrations of mineral dust were recorded in summer compared to winter, with the summer maxima of mineral dust being more evident in the southern and east-

ern European countries. The highest average dust load (5 to 10 $\mu\text{g m}^{-3}$) was determined at eastern and south-western sites (MD13, IT01, ES22, AM01 and ES1778) accounting for 20–40 % of PM₁₀. Intermediate levels (between 2.5 and 5 $\mu\text{g m}^{-3}$) were obtained at GR02, ESCLI, HU02, SK06, FR09, FR22 and IT04, followed by CH02 and DE44 (1.5–2 $\mu\text{g m}^{-3}$, 14–25 % of PM₁₀). At the remaining sites (FR30, SE12, GB48 and IE31), the dust load was <0.5 $\mu\text{g m}^{-3}$ (2–10 % of PM₁₀). In winter, the concentrations of mineral dust were lower (<1–2 $\mu\text{g m}^{-3}$, <5–10 % of PM₁₀) for all sites except for GR02, where an average concentration of 13.5 $\mu\text{g m}^{-3}$ (34 % of PM₁₀) was calculated due to the influence of short-lived, but intense, Saharan dust events.

Mineral dust was attributed to different origins. Saharan dust outbreaks were responsible for increases in mineral dust levels at sites in southern and central Europe in summer, whereas high levels of mineral dust at eastern European sites were attributed to local or regional sources. Results obtained during the EMEP IMPs evidenced that even during NO-SDE the contribution of mineral dust can be significant at southern and eastern European countries, ranging from 10 to 25 % of PM₁₀.

The influence of two African dust outbreaks on the levels and composition of PM₁₀ across Europe was clearly detected. Contribution of mineral dust during SDEs may account for exceedances of the daily limit value established by the European directive on air quality (EC, 2008) at ES1778 and ES22 in summer, and at GR02 in winter. The comparison between the average mineral loads estimated for the whole period and for the non-SDE permitted us to estimate the Saharan dust contribution at each site, ranging from 0.1 to 0.4 $\mu\text{g m}^{-3}$ at the northern sites affected by the SDE, to 0.5–5 $\mu\text{g m}^{-3}$ at the southern sites. It is well known that African dust can significantly contribute to increase PM₁₀ levels in southern Europe (Querol et al., 2009; Pey et al., 2013), although this contribution may significantly vary inter-annually due the occurrence of extreme dust events. Therefore, the number of exceedances due to the contribution of African dust is subject to large variability, and highly dependent on changes in the meteorology. As shown by EEA (2012), exceedances attributable to African dust episodes are usually lower than 20 % of the total exceedances, and account for less than 5 % of the days in the western basin and up to 10 % of the days in the eastern basin. It is interesting to compare these data with PM₁₀ measurements in the Caribbean (Prospero et al., 2014) where the PM₁₀ exceedances of the daily limit value are frequent and driven by African dust, accounting for 10 % of the days on an annual basis.

The impact of the Saharan dust intrusions during the EMEP IMPs resulted in a relative increase of SiO₂ and Al₂O₃ and a relative decrease of CaO, K₂O and MgO. The composition of African dust affecting different regions of Europe was quite homogeneous over the different areas, showing that changes during dust transport were limited for this

study. These results are consistent with those obtained from dust collected in Barbados and Miami during Saharan events, showing a high uniformity of dust composition that was attributed to the mixing of dust originating from different source areas during transport (Prospero and Lamb, 2003; Trapp et al., 2010; Muhs et al., 2014). A higher heterogeneity should be expected in this study given the proximity of the study area to the north of Africa, although a longer time series is necessary to further investigate this.

The inherent complexity of different dust sources and composition can result in significant errors if the PM mineral load is estimated by applying factors to the concentration of a single measured element, as generally proposed by a number of studies. In our case, a higher variability of the dust composition was evident in winter, this being partially attributed to the higher relative contribution of anthropogenic dust at this time of the year. The relative contribution of K was also more pronounced in winter at some central and eastern European sites, probably reflecting a higher impact from biomass combustion.

Finally, the spatial distribution of trace metals and metalloids, and sulfate enabled the identification of specific anthropogenic sources at a regional scale: i.e., shipping emissions in the Mediterranean region (V, Ni and SO₄²⁻), metallurgy (Cr, Ni and Mn) in central and eastern Europe, high-temperature processes (As, Pb and SO₄²⁻) in eastern European countries, and traffic (Cu) at sites affected by emissions from nearby cities.

The Supplement related to this article is available online at doi:10.5194/acp-16-6107-2016-supplement.

Acknowledgements. The present work was supported by the Co-operative Programme for Monitoring and Evaluation of the Long-range Transmission of Air pollutants in Europe (EMEP) under UNECE. We also acknowledge support by the European Union Seventh Framework Programme (FP7/2007-2013) through ACTRIS (grant agreement no. 262254) and by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 654109.

The participation of IDAEA-CSIC was supported by the Spanish Ministry of Economy and Competitiveness and FEDER funds under the project PRISMA (CGL2012-39623-C02-1), by the Generalitat de Catalunya (AGAUR 2015 SGR33 and the DGQA). The authors wish to thank the Norwegian Ministry of foreign affairs for support to the measurements in Armenia and Moldova. The French participation to the campaigns was funded by the French Agency of Environment and Energy Management (ADEME, grants 1262C0022 and 1262C0039). Mines Douai acknowledges support from the CaPPA project which is financed by the French National Research Agency (ANR) through the PIA (Programme d'Investissement d'Avenir) under contract ANR-11-LABX-0005-01, the "Nord-Pas de Calais" Regional Council and the European

Regional Development Fund (ERDF). Participation of the ANDRA is also acknowledged for the measurements at OPE-ANDRA. The authors are also indebted to the fieldwork teams at Auchencorth Moss and Harwell (NERC CEH and Riccardo Energy and Environment Staff). The authors would also like to express our gratitude to the Atmospheric Modelling Laboratory from the Barcelona Supercomputing Centre, the Naval Research Laboratory and the SeaWiFS project (NASA) for the provision of the DREAM, NAAPs aerosol maps and the satellite imagery, respectively. The authors wish to thank D. C. Carslaw and K. Ropkins for providing the Openair software used in this paper (Carslaw and Ropkins, 2012; Carslaw, 2012).

Edited by: J. Surratt

References

- Aas, W., Tsyro, S., Bieber, E., Bergström, R., Ceburnis, D., Eller-
mann, T., Fagerli, H., Frölich, M., Gehrig, R., Makkonen, U.,
Nemitz, E., Otjes, R., Perez, N., Perrino, C., Prévôt, A. S. H.,
Putaud, J.-P., Simpson, D., Spindler, G., Vana, M., and Yttri, K.
E.: Lessons learnt from the first EMEP intensive measurement
periods, *Atmos. Chem. Phys.*, 12, 8073–8094, doi:10.5194/acp-
12-8073-2012, 2012.
- Alastuey, A., Querol, X., Castillo, S., Escudero, M., Avila, A.,
Cuevas, E., Torres, C., Romero, P., Exposito, F., and Garcia,
O.: Characterisation of TSP and PM_{2.5} at Izaña and
Sta. Cruz de Tenerife (Canary Islands, Spain) during a Saharan
Dust Episode (July 2002), *Atmos. Environ.*, 39, 4715–4728,
doi:10.1016/j.atmosenv.2005.04.018, 2005.
- Alemón, E., Herrera, L., Ortiz, E., and Longoria, L. C.: Instrumental
nuclear activation analysis (INAA) characterization of environmental
air filter samples., *Appl. Radiat. Isotopes*, 60, 815–23,
doi:10.1016/j.apradiso.2004.01.004, 2004.
- Alleman, L. Y., Lamaison, L., Perdrix, E., Robache, A., and
Galloo, J.-C.: PM₁₀ metal concentrations and source identification
using positive matrix factorization and wind sectoring in a French
industrial zone, *Atmos. Res.*, 96, 612–625,
doi:10.1016/j.atmosres.2010.02.008, 2010.
- Aluko, O. and Noll, K. E.: Deposition and suspension of large,
airborne particles, *Aerosol Sci. Tech.*, 40, 503–513,
doi:10.1080/02786820600664152, 2006.
- Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A., and
Moreno, T.: Spatial and chemical patterns of PM₁₀ in road dust
deposited in urban environment, *Atmos. Environ.*, 43, 1650–
1659, doi:10.1016/j.atmosenv.2008.12.009, 2009.
- Andreae, M. O.: Soot carbon and excess fine potassium: Long-range
transport of combustion-derived aerosols, *Science*, 220, 1148–
1151, doi:10.1126/science.220.4602.1148, 1983.
- Aymoz, G., Jaffrezo, J.-L., Jacob, V., Colomb, A., and George, Ch.:
Evolution of organic and inorganic components of aerosol during
a Saharan dust episode observed in the French Alps, *Atmos. Chem. Phys.*, 4,
2499–2512, doi:10.5194/acp-4-2499-2004,
2004.
- Bates, T. S., Lamb, B. K., Guenther, A., Dignon, J., and Stoiber,
R. E.: Sulfur emission to the atmosphere from natural sources. *J.
Atmos. Chem.*, 14, 315–337, 1992.
- Bergametti, G., Dutot, A.-L., Buat-Ménard, P., Losno, R., and
Remoudaki, E.: Seasonal variability of the elemental composition
of atmospheric aerosol particles over the northwestern
Mediterranean, *Tellus B*, 41B, 353–361, doi:10.1111/j.1600-
0889.1989.tb00314.x, 1989.
- Carslaw, D. C.: The openair manual – open-source tools for
analysing air pollution data, Manual for version 0.7-0, King’s
College, London, UK, 2012.
- Carslaw, D. C. and Ropkins, K.: Openair – an R package for air
quality data analysis, *Environ. Modell. Softw.*, 27–28, 52–61,
2012.
- Claquin, T., Schulz, M., and Balkanski, Y. J.: Modeling the mineral-
ogy of atmospheric dust sources, *J. Geophys. Res.*, 104, 22243–
22256, 1999.
- Dayan, U., Ziv, B., Shoob, T., and Enzel, Y.: Suspended dust over
southeastern Mediterranean and its relation to atmospheric circulations,
Int. J. Climatol., 28, 915–924, doi:10.1002/joc.1587,
2008.
- Dentener, F. J., Carmichael, G. R., Zhang, Y., Lelieveld, J., and
Crutzen, P. J.: Role of mineral aerosol as a reactive surface in
the global troposphere, *J. Geophys. Res.-Atmos.*, 101, 22869–
22889, 1996.
- Diaz, J., Tobias, A., and Linares, C.: Saharan dust and association
between particulate matter and case-specific mortality: a case–crossover
analysis in Madrid (Spain), *Environ. Health*, 11,
doi:10.1186/1476-069X-11-11, 2012.
- Drever, J. L.: *The Geochemistry of Natural Waters: Surface and
Groundwater Environments*, Prentice Hall, Upper Saddle River,
New Jersey, USA, 1997.
- EC: Directive 2008/50/EC of the European Parliament and of the
council of 21 May 2008 on ambient air quality and cleaner air
for Europe, *Off. J. Eur. Comm.*, L152, 11/6/2008, 1–44, 2008.
- EEA: Particulate matter from natural sources and related reporting
under the EU Air Quality Directive in 2008 and 2009, Technical
report No. 10/2012, ISBN 978-92-9213-325-2, ISSN 1725-2237,
doi:10.2800/55574, 2012.
- Engelstaedter, S., Tegen, I., and Washington, R.: North African dust
emissions and transport, *Earth-Sci. Rev.*, 79, 73–100, 2006.
- Escudero, M., Querol, X., Pey, J., Alastuey, A., Pérez, N., Ferreira,
F., Alonso, S., Rodríguez, S., and Cuevas, E.: A methodology for
the quantification of the net African dust load in air quality
monitoring networks, *Atmos. Environ.*, 41, 5516–5524,
doi:10.1016/j.atmosenv.2007.04.047, 2007.
- Formenti, P., Rajot, J. L., Desboeufs, K., Saïd, F., Grand, N.,
Chevaillier, S., and Schmechtig, C.: Airborne observations of
mineral dust over western Africa in the summer Monsoon
season: spatial and vertical variability of physico-chemical
and optical properties, *Atmos. Chem. Phys.*, 11, 6387–6410,
doi:10.5194/acp-11-6387-2011, 2011.
- Ginoux, P., Prospero, J. M., Gill, T. E., Hsu, N. C., and
Zhao, M.: Global-scale attribution of anthropogenic and natural
dust sources and their emission rates based on MODIS
Deep Blue aerosol products, *Rev. Geophys.*, 50, RG3005,
doi:10.1029/2012RG000388, 2012.
- Harrison, R. M. and Pio, C. A.: Size-differentiated composition
of inorganic atmospheric aerosols of both marine and polluted
continental origin, *Atmos. Environ.*, 17, 1733–1738,
doi:10.1016/0004-6981(83)90180-4, 1983.

- Highwood, E. J. and Ryder, C. L.: Radiative effects of dust, in: *Mineral Dust: A Key Player in the Earth System*, edited by: Knippertz, P. and Stuut, J. W., Springer Science Business Media, Dordrecht, the Netherlands, 2014.
- Huneus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J.-J., Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison in AeroCom phase I, *Atmos. Chem. Phys.*, 11, 7781–7816, doi:10.5194/acp-11-7781-2011, 2011.
- Hwang, H. and Ro, C.-U.: Direct observation of nitrate and sulfate formations from mineral dust and sea-salts using low-Z particle electron probe X-ray microanalysis, *Atmos. Environ.*, 40, 3869–3880, doi:10.1016/j.atmosenv.2006.02.022, 2006.
- IPCC: *Climate Change 2007: The Physical Science Basis*, edited by: Solomon, S., Qin, D., Manning, M., Marquis, M., Averyt, K., Tignor, M. M. B., Miller, H. L., and Chen, Z., Cambridge University Press, Cambridge, UK, 996 pp., 2007.
- IPCC: *Summary for Policymakers*, in: *Climate Change 2013: The Physical Science Basis*, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.
- Journet, E., Balkanski, Y., and Harrison, S. P.: A new data set of soil mineralogy for dust-cycle modeling, *Atmos. Chem. Phys.*, 14, 3801–3816, doi:10.5194/acp-14-3801-2014, 2014.
- Kallos, G., Papadopoulos, A., Katsafados, P., and Nickovic, S.: Transatlantic Saharan dust transport: Model simulation and results, *J. Geophys. Res.*, 111, D09204, doi:10.1029/2005JD006207, 2006.
- Kandler, K., Benker, N., Bundke, U., Cuevas, E., Ebert, M., Knippertz, P., Rodríguez, S., Schütz, L., and Weinbruch, S.: Chemical composition and complex refractive index of Saharan Mineral Dust at Izaña, Tenerife (Spain) derived by electron microscopy, *Atmos. Environ.*, 41, 8058–8074, doi:10.1016/j.atmosenv.2007.06.047, 2007.
- Kandler, K., Schütz, L., Deutscher, C., Ebert, M., Hofmann, H., Jäckel, S., Jaenicke, R., Knippertz, P., Lieke, K., Massling, A., Petzold, A., Schladitz, A., Weinzierl, B., Wiedensohler, A., Zorn, S., and Weinbruch, S.: Size distribution, mass concentration, chemical and mineralogical composition and derived optical parameters of the boundary layer aerosol at Tinfou, Morocco, during SAMUM 2006, *Tellus B*, 61, 32–50, doi:10.1111/j.1600-0889.2008.00385.x, 2009.
- Karanasiou, A., Diapouli, E., Cavalli, F., Eleftheriadis, K., Viana, M., Alastuey, A., Querol, X., and Reche, C.: On the quantification of atmospheric carbonate carbon by thermal/optical analysis protocols, *Atmos. Meas. Tech.*, 4, 2409–2419, doi:10.5194/amt-4-2409-2011, 2011.
- Karanasiou, A., Moreno, N., Moreno, T., Viana, M., de Leeuw, F., and Querol, X.: Health effects from Sahara dust episodes in Europe: literature review and research gaps, *Environ. Int.*, 47, 107–114, 2012.
- Knippertz, P. and Stuut, J. W. (Eds): *Mineral Dust: A Key Player in the Earth System*, Springer Science Business Media, Dordrecht, the Netherlands, 2014.
- Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle, *P. Natl. Acad. Sci. USA*, 108, 1016–1021, 2011.
- Krueger, B. J., Grassian, V. H., Cowin, J. P., and Laskin, A.: Heterogeneous chemistry of individual mineral dust particles from different dust source regions: the importance of particle mineralogy, *Atmos. Environ.*, 38, 6253–6261, doi:10.1016/j.atmosenv.2004.07.010, 2004.
- Lafon, S., Sokolik, I. N., Rajot, J. L., Caquineau, S., and Gaudichet, A.: Characterization of iron oxides in mineral dust aerosols: Implications for light absorption, *J. Geophys. Res.-Atmos.*, 111, D21207, doi:10.1029/2005JD007016, 2006.
- Levin, Z. and Ganor, E.: The effects of desert particles on cloud and rain formation in the eastern Mediterranean, in: *The Impact of Desert Dust Across the Mediterranean*, edited by: Guerzoni, S. and Chester, R., Kluwer Academic Publishers, Dordrecht, the Netherlands, 77–86, 1996.
- Lucarelli, F., Nava, S., Calzolari, G., Chiari, M., Udisti, R., and Marino, F.: Is PIXE still a useful technique for the analysis of atmospheric aerosols? The LABEC experience, *X-Ray Spectrom.*, 40, 162–167, doi:10.1002/xrs.1312, 2011.
- Lucarelli, F., Calzolari, G., Chiari, M., Giannoni, M., Mochi, D., Nava, S., and Carraresi L.: The upgraded external-beam PIXE/PIGE set-up at LABEC for very fast measurements on aerosol samples, *Nucl. Instrum. Meth. B*, 318, 55–59, doi:10.1016/j.nimb.2013.05.099, 2014.
- Maenhaut, W.: Present role of PIXE in atmospheric aerosol research, *Nucl. Instrum. Meth. B*, 363, 86–91, doi:10.1016/j.nimb.2015.07.043, 2015.
- Mallone, S., Stafoggia, M., Faustini, A., Gobbi, G. P., Marconi, A., and Forastiere, F.: Saharan dust and associations between particulate matter and daily mortality in Rome, Italy, *Environ. Health Perspect.*, 119, 1409–1414, 2011.
- McMeeking, G. R., Kreidenweis, S. M., Baker, S., Carrico, C. M., Chow, J. C., Collett, J. L., Hao, W. M., Holden, A. S., Kirchstetter, T. W., Malm, W. C., Moosmüller, H., Sullivan, A. P., and Cyle E., W.: Emissions of trace gases and aerosols during the open combustion of biomass in the laboratory, *J. Geophys. Res.-Atmos.*, 114, D19210, doi:10.1029/2009JD011836, 2009.
- Miller, R. L., Knippertz, P., García-Pando, C. P., Perlwitz, J. P., and Tegen, I.: Impact of dust radiative forcing upon climate, in: *Mineral Dust: A Key Player in the Earth System*, edited by: Knippertz, P. and Stuut, J. W., Springer Science Business Media, Dordrecht, the Netherlands, 2014.
- Mitsakou, C., Kallos, G., Papantoniou, N., Spyrou, C., Solomos, S., Astitha, M., and Housiadas, C.: Saharan dust levels in Greece and received inhalation doses, *Atmos. Chem. Phys.*, 8, 7181–7192, doi:10.5194/acp-8-7181-2008, 2008.
- Moreno, T., Querol, X., Castillo, S., Alastuey, A., Cuevas, E., Herrmann, L., Mounkaila, M., Elvira, J., and Gibbons, W.: Geochemical variations in aeolian mineral particles from the Sahara-Sahel Dust Corridor, *Chemosphere*, 65, 261–270, doi:10.1016/j.chemosphere.2006.02.052, 2006.
- Moreno, T., Amato, F., Querol, X., Alastuey, A., and Gibbons, W.: Trace element fractionation processes in resuspended mineral

- aerosols extracted from Australian continental surface materials, *Aust. J. Soil Res.*, 46, 128–140, doi:10.1071/SR07121, 2008.
- Morman, S. A. and Plumlee, G. S.: Dust and human health, in: *Mineral Dust: A Key Player in the Earth System*, edited by: Knippertz, P. and Stuut, J. W., Springer Science Business Media, Dordrecht, the Netherlands, 385–409, 2014.
- Moulin, C., Lambert, C. E., Dayan, U., Masson, V., Ramonet, M., Bousquet, P., Legrand, M., Balkanski, Y. J., Guelle, W., Marticorena, B., Bergametti, G., and Dulac, F.: Satellite climatology of African dust transport in the Mediterranean atmosphere, *J. Geophys. Res.*, 103, 13137, doi:10.1029/98JD00171, 1998.
- Muhs, D. R., Prospero, J. M., Baddock, M. C., Gill, T. E., Muhs, D. R., Prospero, J. M., Baddock, M. C., and Gill, T. E.: Identifying Sources of Aeolian Mineral Dust: Present and Past, in: *Mineral Dust: A Key Player in the Earth System*, edited by: Knippertz, P. and Stuut, J. W., Springer Science Business Media, Dordrecht, the Netherlands, 2014.
- Nava, S., Becagli, S., Calzolari, G., Chiari, M., Lucarelli, F., Prati, P., Traversi, R., Udisti, R., Valli, G., and Vecchi, R.: Saharan dust impact in central Italy: An overview on three years elemental data records, *Atmos. Environ.*, 60, 444–452, doi:10.1016/j.atmosenv.2012.06.064, 2012.
- Nenes, A., Murray, B., and Bougiatioti, A.: Mineral Dust and its Microphysical Interactions with Clouds, in: *Mineral Dust: A Key Player in the Earth System*, edited by: Knippertz, P. and Stuut, J. W., Springer Science Business Media, Dordrecht, the Netherlands, 2014.
- Nozaki, Y.: A Fresh Look at Element Distribution in the North Pacific, *Eos Trans. AGU*, 78, 221–221, doi:10.1029/97EO00148, 1997.
- Nriagu, J. O. and Pacyna, J. M.: Quantitative assessment of worldwide contamination of air, water and soils by trace metals, *Nature*, 333, 134–139, doi:10.1038/333134a0, 1988.
- Pacyna J. M.: Atmospheric trace elements from natural and anthropogenic sources, in: *Toxic Metals in the Atmosphere*, edited by: Nriagu, J. O. and Davidson, C. I., Wiley, New York, USA, 33–50, 1986.
- Pandolfi, M., Gonzalez-Castanedo, Y., Alastuey, A., de la Rosa, J. D., Mantilla, E., de la Campa, A. S., Querol, X., Pey, J., Amato, F., and Moreno, T.: Source apportionment of PM₁₀ and PM_{2.5} at multiple sites in the strait of Gibraltar by PMF: impact of shipping emissions, *Environ. Sci. Pollut. R.*, 18, 260–169, doi:10.1007/s11356-010-0373-4, 2011.
- Pérez, L., Tobias, A., Querol, X., Künzli, N., Pey, J., Alastuey, A., Viana, M., Valero, N., González-Cabré, M., and Sunyer, J.: Coarse Particles From Saharan Dust and Daily Mortality, *Epidemiology*, 19, 800–807, doi:10.1097/EDE.0b013e31818131cf, 2008.
- Pérez, L., Tobías, A., Pey, J., Pérez, N., Alastuey, A., Sunyer, J., and Querol, X.: Effects of Local and Saharan Particles on Cardiovascular Disease Mortality, *Epidemiology*, 23, 768–769, doi:10.1097/EDE.0b013e3182625d0d, 2012.
- Pey, J., Querol, X., Alastuey, A., Forastiere, F., and Stafoggia, M.: African dust outbreaks over the Mediterranean Basin during 2001–2011: PM₁₀ concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology, *Atmos. Chem. Phys.*, 13, 1395–1410, doi:10.5194/acp-13-1395-2013, 2013.
- Prospero, J. M. and Lamb, P. J.: African droughts and dust transport to the Caribbean: climate change implications, *Science*, 302, 1024–1027, 2003.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40, 1002, doi:10.1029/2000RG000095, 2002.
- Prospero, J. M., Collard, F.-X., Molinié, J., and Jeannot, A.: Characterizing the annual cycle of African dust transport to the Caribbean Basin and South America and its impact on the environment and air quality, *Global Biogeochem. Cy.*, 28, 757–773, doi:10.1002/2013GB004802, 2014.
- Putaud, J. P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W., Cyrys, J., Flentje, H., Fuzzi, S., Gehrig, R., Hansson, H. C., Harrison, R. M., Herrmann, H., Hitzenberger, R., Hüglin, C., Jones, A. M., Kasper-Giebl, A., Kiss, G., Kousa, A., Kuhlbusch, T. A. J., Löschau, G., Maenhaut, W., Molnar, A., Moreno, T., Pekkanen, J., Perrino, C., Pitz, M., Puxbaum, H., Querol, X., Rodriguez, S., Salma, I., Schwarz, J., Smolik, J., Schneider, J., Spindler, G., ten Brink, H., Tursic, J., Viana, M., Wiedensohler, A., and Raes, F.: A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.*, 44, 1308–1320, doi:10.1016/j.atmosenv.2009.12.011, 2010.
- Querol, X., Alastuey, A., Puigercus, J., Mantilla, E., Ruiz, C. R., Lopez-Soler, A., Plana, F., and Juan, R.: Seasonal evolution of suspended particles around a large coal-fired power station: Chemical characterization, *Atmos. Environ.*, 32, 719–731, doi:10.1016/S1352-2310(97)00340-3, 1998.
- Querol, X., Alastuey, A., Rodriguez, S., Plana, F., Ruiz, C. R., Cots, N., Massagué, G., and Puig, O.: PM₁₀ and PM_{2.5} source apportionment in the Barcelona Metropolitan area, Catalonia, Spain, *Atmos. Environ.*, 35, 6407–6419, doi:10.1016/S1352-2310(01)00361-2, 2001.
- Querol, X., Alastuey, A., Rodríguez, S., Viana, M. M., Artíñano, B., Salvador, P., Mantilla, E., Dos Santos, S. G., Patier, R. F., De La Rosa, J., De La Campa, A. S., Menéndez, M., and Gil, J. J.: Levels of particulate matter in rural, urban and industrial sites in Spain, *Sci. Total Environ.*, 334–335, 359–376, doi:10.1016/j.scitotenv.2004.04.036, 2004.
- Querol, X., Viana, M., Alastuey, A., Amato, F., Moreno, T., Castillo, S., Pey, J., de la Rosa, J., Sánchez de la Campa, A., Artíñano, B., Salvador, P., García Dos Santos, S., Fernández-Patier, R., Moreno-Grau, S., Negral, L., Minguillón, M. C., Monfort, E., Gil, J. I., Inza, A., Ortega, L. A., Santamaría, J. M., and Zabalza, J.: Source origin of trace elements in PM from regional background, urban and industrial sites of Spain, *Atmos. Environ.*, 41, 7219–7231, doi:10.1016/j.atmosenv.2007.05.022, 2007.
- Querol, X., Pey, J., Pandolfi, M., Alastuey, A., Cusack, M., Pérez, N., Moreno, T., Viana, M., Mihalopoulos, N., Kallos, G., and Kleanthous, S.: African dust contributions to mean ambient PM₁₀ mass-levels across the Mediterranean Basin, *Atmos. Environ.*, 43, 4266–4277, doi:10.1016/j.atmosenv.2009.06.013, 2009.
- Rodríguez, S., Querol, X., Alastuey, A., Kallos, G., and Kakaliagou, O.: Saharan dust contributions to PM₁₀ and TSP levels in Southern and Eastern Spain, *Atmos. Environ.*, 35, 2433–2447, doi:10.1016/S1352-2310(00)00496-9, 2001.

- Rodríguez, S., Alastuey, A., Alonso-Pérez, S., Querol, X., Cuevas, E., Abreu-Afonso, J., Viana, M., Pérez, N., Pandolfi, M., and de la Rosa, J.: Transport of desert dust mixed with North African industrial pollutants in the subtropical Saharan Air Layer, *Atmos. Chem. Phys.*, 11, 6663–6685, doi:10.5194/acp-11-6663-2011, 2011.
- Rodríguez, S., Alastuey, A., and Querol, X.: A review of methods for long term in situ characterization of aerosol dust, *Aeolian Res.*, 6, 55–74, doi:10.1016/j.aeolia.2012.07.004, 2012.
- Schauer, J. J., Lough, G. C., Shafer, M. M., Christensen, W. F., Arndt, M. F., DeMinter, J. T., and Park, J.-S.: Characterization of Metals Emitted from Motor Vehicles, Research Report 133, Health Effects Institute, 1–76; discussion 77–88, Boston, MA, USA, 2006.
- Scheuvens, D., Kandler, K., Küpper, M., Lieke, K., Zorn, S. R., Ebert, M., Schütz, L., and Weinbruch, S.: Individual-particle analysis of airborne dust samples collected over Morocco in 2006 during SAMUM 1, *Tellus B*, 63, 512–530, doi:10.1111/j.1600-0889.2011.00554.x, 2011.
- Scheuvens, D., Schütz, L., Kandler, K., Ebert, M., and Weinbruch, S.: Bulk composition of northern African dust and its source sediments – A compilation, *Earth-Sci. Rev.*, 116, 170–194, doi:10.1016/j.earscirev.2012.08.005, 2013.
- Stafoggia, M., Samoli, E., Alessandrini, E., Cadum, E., Ostro, B., Berti, G., Faustini, A., Jacquemin, B., Linares, C., Pascal, M., Randi, G., Ranzì, A., Stivanello, E., and Forastiere, F.: Short-term associations between fine and coarse particulate matter and hospitalizations in Southern Europe: Results from the MED-PARTICLES project, *Environ. Health Perspect.*, 121, 1026–1033, doi:10.1289/ehp.1206151, 2013.
- Tegen, I., Werner, M., Harrison, S. P., and Kohfeld, K. E.: Relative importance of climate and land use in determining present and future global soil dust emission, *Geophys. Res. Lett.*, 31, L05105, doi:10.1029/2003GL019216, 2004.
- Tørseth, K., Aas, W., Breivik, K., Fjæraa, A. M., Fiebig, M., Hjellbrekke, A. G., Lund Myhre, C., Solberg, S., and Yttri, K. E.: Introduction to the European Monitoring and Evaluation Programme (EMEP) and observed atmospheric composition change during 1972–2009, *Atmos. Chem. Phys.*, 12, 5447–5481, doi:10.5194/acp-12-5447-2012, 2012.
- Trapp, J. M., Millero, F. J., and Prospero, J. M.: Temporal variability of the elemental composition of African dust measured in trade wind aerosols at Barbados and Miami, *Mar. Chem.*, 120, 71–82, doi:10.1016/j.marchem.2008.10.004, 2010.
- UNECE: EMEP monitoring strategy for 2010–2019, United Nations Economic Commissions for Europe, Geneva, Switzerland, ECE/EB.AIR/GE.1/2009/15, available at: <http://www.unece.org/env/documents/2009/EB/ge1/ece.eb.air.ge.1.2009.15.e.pdf> (last access: December 2015), 2009.
- Viana, M., Kuhlbusch, T. A. J., Querol, X., Alastuey, A., Harrison, R. M., Hopke, P. K., Winiwarter, W., Vallius, M., Szidat, S., Prévôt, A. S. H., Hueglin, C., Bloemen, H., Wählin, P., Vecchi, R., Miranda, A. I., Kasper-Giebl, A., Maenhaut, W., and Hitzenberger, R.: Source apportionment of particulate matter in Europe: A review of methods and results, *J. Aerosol Sci.*, 39, 827–849, doi:10.1016/j.jaerosci.2008.05.007, 2008.
- WMO: WMO/GAW Aerosol Measurement Procedures Guidelines and Recommendations, GAW Report No. 153 (WMO TD No. 1178), September 2003, World Meteorological Organization, Scientific Advisory Group Aerosols, Geneva, Switzerland, 67 pp., 2003.
- WMO: WMO/GAW Strategic Plan: 2008–2015 – A Contribution to the Implementation of the WMO Strategic Plan: 2008–2011, GAW Report No. 172 (WMO TD No. 1384), July 2007, World Meteorological Organization, Geneva, Switzerland, 108 pp., 2007.
- Zender, C. S., Miller, R. L. R. L., and Tegen, I.: Quantifying mineral dust mass budgets: Terminology, constraints, and current estimates, *Eos T. Am. Geophys. Un.*, 85, 509–512, doi:10.1029/2004EO480002, 2004.
- Zhao, C., Liu, X., Leung, L. R., Johnson, B., McFarlane, S. A., Gustafson Jr., W. I., Fast, J. D., and Easter, R.: The spatial distribution of mineral dust and its shortwave radiative forcing over North Africa: modeling sensitivities to dust emissions and aerosol size treatments, *Atmos. Chem. Phys.*, 10, 8821–8838, doi:10.5194/acp-10-8821-2010, 2010.